

***Camissonia nevadensis*** (Kellogg [1862]) Raven**NEVADA SUNCUP****FAMILY:** *Onagraceae*, the evening-primrose family.**SYNONYMS:** *Oenothera nevadensis*  
*Sphaerostigma tortuosa***STATUS:****Heritage Program WATCH LIST, ranks: G3 S3**

USFWS/ESA: none. STATE OF NEVADA: none. BLM: none. USFS: none. NNNPS: none.

**POPULATION CENSUS:** 7 occurrences mapped; total estimated individuals 506+, total estimated area unknown.**TREND:** unknown.**IMPACTS AND MAJOR THREATS:** No summary available (see references).**INVENTORY EFFORT:** Systematic surveys have not yet been conducted. Most recent entered survey 1989, average year of last survey 1980. **Years since last entered survey** (percent of mapped records at various survey ages): **11-20 yrs:** 14.3%; **21-30 yrs:** 85.7%.**LAND MANAGEMENT:** U. S. Bureau of Land Management.**RANGE:** Carson City, Churchill, Douglas, Lyon, Pershing, Storey, and Washoe counties, Nevada. **Nevada endemic.** Maximum **range dimension 128.3 km (79.8 mi)** excluding most disjunct record.**ELEVATIONS RECORDED:** 4050-5250 feet (1234-1600 meters).**HABITAT:** Open, sandy, gravelly, or clay slopes and flats in the salt-desert, shadscale, and lower sagebrush zones.**PHENOLOGY:** flowering spring. Range of most frequent **survey months:** April-May.**LIFE-FORM AND HABIT:** small annual.**DESCRIPTION:** not available (see references).**PHOTOGRAPHS:** Nevada Natural Heritage Program images web page (1998-present).**ILLUSTRATIONS:** Cronquist et al. (1997), Mozingo and Williams (1980).**SPECIFIC REFERENCES:**

Kellogg, A. 1862. [Untitled]. Proceedings of the California Academy of Sciences 2: 179, 185-201, 224-232.